

A Review On The Role of Homoeopathy In The Management of Hyperuricemia



Dr. Aditi B. Bhise

Abstract-Hyperuricemia is a metabolic condition characterized by elevated levels of uric acid in the blood, which may lead to various complications such as gout, renal disorders, and joint inflammation. The increasing prevalence of hyperuricemia worldwide has become a significant health concern due to changing lifestyle patterns, dietary habits, and associated metabolic disorders. Conventional treatment primarily focuses on lowering uric acid levels through pharmacological agents; however, long-term use of such medications may be associated with adverse effects and recurrence of symptoms.

Homoeopathy, based on the principle of Similia Similibus Curentur (let like be cured by like), offers a holistic and individualized approach to the management of metabolic disorders including hyperuricemia. Homoeopathic medicines aim not only to reduce uric acid levels but also to improve the patient's overall constitution, thereby addressing the underlying predisposition of the disease. Remedies such as Colchicum autumnale, Benzoic acid, Ledum palustre, Lycopodium clavatum, and Urtica urens are frequently indicated in cases presenting with symptoms related to hyperuricemia and gout.

This review aims to analyze and compile the available literature regarding the role of homoeopathy in the management of hyperuricemia. Various classical homoeopathic texts, repertories, and research articles have been studied to understand the therapeutic indications of commonly used remedies and their clinical relevance. The study highlights the scope of homoeopathic treatment in reducing symptoms, improving quality of life, and preventing complications associated with hyperuricemia.

Keywords:Hyperuricemia, Serum Uric Acid, Metabolic disorder, Homoeopathy.

Introduction: Hyperuricemia is a metabolic condition characterized by an elevated level of uric acid in the blood, usually exceeding 7 mg/dL in men and 6 mg/dL in women. Uric acid is the final product of purine metabolism in the human body, and it is normally eliminated through the kidneys and, to a lesser extent, through the gastrointestinal tract. When the production of uric acid increases or

its excretion decreases, uric acid accumulates in the bloodstream, leading to hyperuricemia. This condition may remain asymptomatic for long periods but can eventually lead to various clinical manifestations, most notably gout, renal calculi, and chronic kidney disease.

The prevalence of hyperuricemia has been increasing globally due to changes in lifestyle, dietary habits, and the rising incidence of metabolic disorders such as obesity, hypertension, diabetes mellitus, and dyslipidemia. High intake of purine-rich foods, alcohol consumption, sedentary lifestyle, and certain medications are recognized contributing factors. Persistent hyperuricemia may lead to deposition of monosodium urate crystals in joints and tissues, resulting in inflammatory conditions like gouty arthritis and other systemic complications.

Homoeopathy, based on the principles of "Similia Similibus Curentur" and individualization, offers a holistic approach in the management of metabolic disorders including hyperuricemia. Homoeopathic treatment aims not only at reducing the serum uric acid levels but also at correcting the underlying constitutional imbalance, thereby improving the patient's overall health and preventing recurrence of symptoms. Several homoeopathic remedies such as Colchicum, Ledum palustre, Benzoic acid, Lycopodium, and Urtica urens have been traditionally used in conditions associated with elevated uric acid levels and gouty manifestations.

Therefore, exploring the role of homoeopathy in the management of hyperuricemia is important to understand its potential in providing a safe, effective, and holistic therapeutic approach. This review aims to analyze the available literature regarding the role of homoeopathic medicines in managing hyperuricemia and its associated clinical conditions, thereby highlighting the scope of homoeopathy in this increasingly prevalent metabolic disorder.

Definition: Hyperuricemia is defined as an abnormally elevated level of uric acid in the blood resulting from increased production or decreased excretion of uric acid. Clinically, hyperuricemia is generally considered when the serum uric acid concentration exceeds 7 mg/dL in adult males and 6 mg/dL in adult females, which is above the normal physiological solubility limit of uric acid in plasma.


Epidemiology: Globally, the prevalence of hyperuricemia varies between 10% and 25% in the general population, depending on the population studied, dietary patterns, and genetic predisposition. Developed countries have reported higher prevalence rates due to increased consumption of purine-rich foods, alcohol intake, obesity, and sedentary lifestyles. Studies have also shown that hyperuricemia is more common in men than in women, particularly in middle-aged and elderly populations. In women, the prevalence increases after menopause because of reduced estrogen levels, which normally help in the excretion of uric acid.

In India, the prevalence of hyperuricemia has been increasing in recent years due to rapid urbanization, changes in diet, and increasing rates of obesity and metabolic disorders. Various epidemiological studies suggest that the prevalence in the Indian population ranges from 9% to 25%, with higher rates reported in urban populations compared to rural areas. The condition is frequently associated with comorbidities such as hypertension, diabetes mellitus, obesity, and chronic kidney disease.

Causes: Hyperuricemia occurs when there is an imbalance between the production and excretion of uric acid in the body. It may result either from increased production of uric acid, decreased renal excretion, or a combination of both. The causes of hyperuricemia can be broadly classified into the following categories:

•Increased Production of Uric Acid - Hyperuricemia may occur due to excessive formation of uric acid in the body. This can result from increased breakdown of purines or excessive intake of purine-rich foods. Diets rich in red meat, organ meat, seafood, and alcoholic beverages can increase uric acid production. Certain metabolic conditions and diseases that involve rapid cell turnover, such as leukemia, lymphoma, and psoriasis, can also lead to increased purine metabolism and elevated uric acid levels.

•Decreased Excretion of Uric Acid - The kidneys play a major role in eliminating uric acid from the body. When renal excretion is impaired, uric acid accumulates in the bloodstream, leading to hyperuricemia. Chronic kidney disease, dehydration, and certain medications such as diuretics, low-dose aspirin, and cyclosporine may



reduce uric acid excretion.

•Dietary Factors - Excessive consumption of purine-rich foods such as red meat, shellfish, and organ meats, as well as beverages high in fructose (soft drinks and fruit juices), can increase uric acid levels. Alcohol intake, particularly beer and spirits, is also strongly associated with hyperuricemia because it increases uric acid production and decreases its excretion.

•Genetic Factors- Some individuals have a hereditary tendency to develop hyperuricemia due to inherited abnormalities in purine metabolism or renal uric acid transport. Genetic predisposition plays an important role in the development of hyperuricemia and gout.

•Associated Medical Conditions - Hyperuricemia is commonly associated with several metabolic and systemic disorders such as obesity, hypertension, diabetes mellitus, metabolic syndrome, hypothyroidism, and chronic kidney disease. These conditions may interfere with normal uric acid metabolism and excretion.

•Drug-Induced Hyperuricemia - Certain medications can increase uric acid levels by either increasing production or reducing excretion. These include thiazide and loop diuretics, low-dose aspirin, pyrazinamide, ethambutol, cyclosporine, and some chemotherapeutic drugs.


Pathophysiology:Hyperuricemia results from an imbalance between the production and excretion of uric acid, the final product of purine metabolism in humans. Purines are derived from dietary sources, cellular turnover, and de novo synthesis in the body.

These purines are metabolized in the liver through a series of enzymatic reactions, ultimately forming uric acid. The enzyme xanthine oxidase plays a key role in converting hypoxanthine to xanthine and xanthine to uric acid.

Under normal physiological conditions, uric acid is dissolved in the blood and eliminated mainly through the kidneys (approximately two-thirds) and partly through the gastrointestinal tract. The kidneys filter uric acid at the glomerulus, after which a portion is reabsorbed and secreted in the renal tubules. When this balance between production and elimination is disturbed, uric acid accumulates in the bloodstream, leading to hyperuricemia.

Hyperuricemia may occur due to overproduction of uric acid, reduced renal excretion, or a combination of both. Overproduction may occur due to increased purine intake, increased breakdown of nucleic acids during rapid cell turnover (as seen in malignancies or psoriasis), or enzymatic defects in purine metabolism. However, in most individuals, hyperuricemia occurs primarily due to decreased renal excretion of uric acid.

Clinical Features:Hyperuricemia often remains asymptomatic for many years and is frequently detected incidentally during routine blood investigations. However, when the level of uric acid rises significantly or persists for a long period, it may lead to various clinical manifestations due to the deposition of monosodium urate crystals in joints, kidneys, and other tissues.



One of the most common clinical manifestations of hyperuricemia is gout, an inflammatory condition characterized by sudden onset of severe joint pain, swelling, redness, and tenderness. The first metatarsophalangeal joint (great toe) is most commonly affected, although other joints such as the ankle, knee, wrist, and fingers may also be involved. The pain usually appears suddenly, often at night, and may be accompanied by warmth and restricted movement of the affected joint.

Another important manifestation is the formation of tophi, which are deposits of monosodium urate crystals in soft tissues. Tophi may develop in chronic cases and are commonly seen around the joints, ears, elbows, fingers, and Achilles tendon. These nodular swellings can cause joint deformity and limitation of movement if left untreated.

Hyperuricemia may also affect the renal system. Uric acid crystals may accumulate in the kidneys leading to uric acid nephrolithiasis (kidney stones), which presents with symptoms such as severe flank pain, hematuria, nausea, and vomiting. Persistent deposition of urate crystals in the kidneys may also result in urate nephropathy, which can gradually impair renal function.

Diagnosis: The diagnosis of hyperuricemia is primarily based on laboratory investigations, clinical evaluation, and identification of associated conditions. It is often detected during routine blood tests or during the evaluation of symptoms related to gout, kidney stones, or metabolic disorders.

The most important diagnostic test is the measurement of serum uric acid levels. Hyperuricemia is generally diagnosed when the serum uric acid level exceeds 7 mg/dL in adult males and 6 mg/dL in adult females. This test helps in identifying elevated uric acid levels even in asymptomatic individuals.

In addition to serum uric acid estimation, urine analysis may be performed to measure the amount of uric acid excreted in the urine. A 24-hour urine uric acid test helps determine whether hyperuricemia is caused by increased production or decreased excretion of uric acid.

Synovial fluid examination may be carried out in patients presenting with joint pain or suspected gout. The presence of monosodium urate crystals in

synovial fluid under polarized light microscopy confirms the diagnosis of gout associated with hyperuricemia.

Imaging studies such as X-ray, ultrasound, or CT scan may be used to detect joint damage, urate crystal deposits, or kidney stones caused by long-standing hyperuricemia. Ultrasound may reveal features such as the "double contour sign," indicating urate crystal deposition on cartilage surfaces.

Additional laboratory tests may also be performed to evaluate associated conditions and complications. These include renal function tests (serum creatinine and blood urea nitrogen), blood glucose levels, lipid profile, and complete blood count. These investigations help assess underlying metabolic disorders and kidney involvement.

Homeopathic Approach: Homeopathy offers a holistic and individualized approach in the management of hyperuricemia. According to homeopathic principles, diseases arise due to disturbances in the vital force, and treatment is directed toward restoring balance in the body rather than merely suppressing symptoms. In hyperuricemia, homeopathic management focuses on correcting the underlying constitutional imbalance, improving metabolism, and enhancing the body's natural ability to eliminate excess uric acid.

The homeopathic approach emphasizes individualization of the patient rather than treating the disease alone. Each patient is evaluated based on their totality of symptoms, including physical complaints, mental and emotional characteristics, modalities, and personal constitution. Based on this comprehensive case-taking, a suitable constitutional remedy is selected that corresponds to the patient's symptom picture.

Homeopathy also considers the miasmatic background of the patient, which may play an important role in chronic metabolic disorders such as hyperuricemia. Remedies are selected according to the dominant miasmatic influence, helping to address the deeper constitutional predisposition and reduce the tendency for recurrence.

Homeopathic Therapeutics:

•**Urtica urens** - This remedy is often indicated in rheumatic complaints affecting the upper limbs, particularly when the deltoid muscles are involved. Patients may experience pain in the right deltoid muscle accompanied by cramp-like sensations that tend to aggravate in the evening. There may also be stiffness and soreness on the inner side of the left knee, along with rheumatic pains affecting both ankles. In some cases, the patient develops rheumatic stiffness and pain in the right wrist, which may later extend to the left arm, wrist, and fingers. Skin symptoms such as raised, red, and intensely itching blisters may appear on the hands and fingers.

•**Colchicum autumnale** - This medicine is particularly suited to individuals with a rheumatic

or gouty constitution. The rheumatic pains tend to shift from one joint to another and are usually aggravated by movement but relieved by cold applications, even though the patient may feel chilly. The complaints are often worsened in cold and damp weather. Weakness of the lower limbs may become marked, sometimes causing the knees to knock together while walking. The great toe may become inflamed and extremely sensitive, with sharp, stitching pains that are aggravated by the slightest touch.

•**Ocimum canum** - This remedy is useful in cases marked by intense colicky pain accompanied by severe vomiting. The patient may twist and writhe in pain, crying out in distress. It is frequently indicated in right-sided renal calculi, where the pain is severe and associated with persistent vomiting. The urine may appear reddish with brick-red sediment following an attack. Renal colic may also be accompanied by numbness in the right thigh. Additionally, there may be pain and swelling in the right ankle joint, which is often more pronounced in the morning.

•**Benzoic Acid** - This remedy is particularly suited to individuals with a gouty and rheumatic constitution, especially when the condition develops in patients with a history of gonorrhoeal or syphilitic infection. The characteristic symptoms include tearing and stitching pains in the large joints, particularly the big toe, accompanied by redness and swelling. These complaints are usually worse at night. A marked feature of this remedy is that the pain frequently shifts from one location to another. The urine and stool possess a strong, offensive, and penetrating odor. The urine is usually dark in color, though at times it may alternate with lighter urine that is passed in large quantities.

•**Ledum Palustre** - This remedy is well indicated in persons with a rheumatic or gouty diathesis, especially those whose constitutions have been weakened by excessive alcohol consumption. The rheumatic symptoms typically begin in the lower limbs and gradually ascend upward. It is particularly useful in patients whose condition has become weak or asthenic after excessive use of colchicum. There may be persistent swelling of the feet, and the soles are painful while walking, as if filled with blood. Patients often experience heat in the hands and feet in the evening, accompanied by prolonged warm perspiration of the palms and soles. Numbness and a "gone to sleep" sensation in the limbs are also commonly observed.

•**Pulsatilla** - This remedy is commonly indicated when joints such as the ankles and the small joints of the hands and feet are affected. There may be swelling of the knee and ankle joints, along with swelling of the fingers and toes, accompanied by drawing and tensive pains. Patients may complain of simple aching pains in the limbs and joints, which may compel them to stretch their body for relief. These symptoms are often associated with a

general sensation of heat in the body and thirst in the morning while in bed.

•**Kali Carbonicum** - The guiding symptoms of this remedy include backache, marked weakness, and profuse perspiration. It is useful in cases of arthritis and deforming arthritis, particularly when the small joints are involved. Patients often experience rheumatic pains that worsen at night. The joint pains are usually sharp and aching, affecting both joints and tendons. Spasmodic contractions of the fingers and toes may also be present. The knees become painful, especially when ascending or descending stairs.

•**Lycopodium** - This remedy is frequently indicated in chronic gouty conditions. Patients experience drawing and tearing pains in the limbs, especially during rest or at night. Tearing pains may affect the shoulder and elbow joints, and the finger joints may become distorted with chalky deposits. There may also be pain in the heels while walking, as if stepping on a small pebble. Numbness in the hands and feet and cracking of joints during movement or stretching are also noted. Lycopodium also has marked action on the urinary system, with pain in the left kidney and ureter. The urine may appear turbid and milky with offensive sediment, accompanied by dull pressing pain in the bladder and abdominal region. It promotes the formation and deposition of lithic acid, leading to discomfort in the kidneys and bladder.

•**Berberis vulgaris** - This remedy is commonly indicated in elderly individuals with a gouty constitution who appear pale, weak, and sensitive to cold. Patients often experience sharp, fleeting pains in the finger joints where uric acid deposits tend to occur, accompanied by numbness and stiffness. There may be marked soreness and pressure in the renal and lumbar regions. Gouty nodules may develop in the auricles of the ears, producing intense pain described as tearing or stinging in nature, which becomes worse on deep pressure. The pain may radiate down the back and extend into the pelvic region along the course of the uterus. Symptoms frequently shift from one joint to another, and the back often feels stiff and numb.

•**Bryonia alba** - This remedy is useful in cases where the joints, particularly the knees, become stiff, swollen, red, and hot with sharp stitching or tearing pains that are aggravated by the slightest movement. There may also be painful stiffness in the nape of the neck along with stitching pain and rigidity in the lumbar region. Symptoms often arise after sudden changes in weather. Joint and limb pains are generally relieved by warmth. Swelling and heat may be present in the feet, ankles, or knees, especially following injury, and the affected parts become painful and worse with motion.

Conclusion: Hyperuricemia is a common metabolic disorder characterized by elevated levels of uric acid in the blood and is often associated with conditions such as gout, renal calculi, and metabolic syndrome. The increasing prevalence of

hyperuricemia due to sedentary lifestyle, dietary habits, and associated systemic diseases highlights the need for effective and safe management strategies. Early diagnosis and appropriate treatment are essential to prevent long-term complications affecting the joints, kidneys, and cardiovascular system.

Homeopathy provides a holistic and individualized approach in the management of hyperuricemia. By considering the totality of symptoms, constitutional characteristics, and miasmatic background of the patient, homeopathic treatment aims to restore the balance of the vital force and improve the body's natural ability to regulate uric acid metabolism. Various homeopathic remedies such as Colchicum, Ledum palustre, Benzoic acid, Urtica urens, and Lycopodium have been found beneficial in managing symptoms associated with elevated uric acid levels and gouty conditions.

- References:**
1. Billa G, Dargad R, Mehta A. Prevalence of Hyperuricemia in Indian Subjects attending Hyperuricemia Screening Programs-A Retrospective Study. The Journal of the Association of Physicians of India. 2018 Apr 1;66(4):43-6.
 2. Ch Pavan Bindu, Mohanty Shruti, Comparative and correlation study of serum uric acid among nondiabetic and diabetic range of fasting blood glucose. International journal of Pharmacy and Biological Sciences; Oct-Dec 2015 ;5(4):155-158.
 3. Satyanarayana U, Chakrepani U, Biochemistry .3rd ed. Kolkatha: Books and Allied ;2004.
 4. K V Krishna Das, Textbook of Medicine ,5th edition p. 729-731.
 5. Harrison T, Faucia. Harrison's principles of internal medicine. New York: McGraw -Hill Medical;2008, pp2445-2446.
 6. Walker B, Colledge N, Ralston S, Penman I, Davidson's principles and practice of medicine .22 ed. 2014.
 7. Das K, Text book of medicine .5th ed. Jaypee Brothers medical publishers;2008
 8. Grassi W, De Angelis R, Clinical features of gout. Rheumatism. 2011;238-45.
 9. Mondal TC. Spirit of organon part. New Delhi. B. Jain Publishers Pvt Ltd; 2008. p.36-38.
 10. Nelson DL, Lehninger AL, Cox M Michel. Lehninger principles of biochemistry. Macmillan; NH Freeman & Company, New York, 4th edn, 2005, p.873-876.

Dr. Aditi B. Bhise, BHMS, PG Scholar,
Department of Practice of Medicine,
Motiwala (National) Homeopathic Medical
College and Hospital, Nashik.

Under The Guidance of
Dr. Ajay Valke, MD (Hom.),
HOD and Professor,

Department of Practice of Medicine,
Motiwala (National) Homeopathic Medical
College and Hospital, Nashik