



PRO LIFE ERA
HOMEOPATHY JOURNAL



Dr. Qureshi
Rukhsar Rajik

Efficacy of Homoeopathic Medicine in The Management Of Nocturnal Enuresis in Pediatric Age Group 4-10 Years

Abstract-Background-

Nocturnal enuresis, commonly known as bed wetting, is a frequent pediatric problem characterized by involuntary passage of urine during sleep. Although it is often regarded as a developmental condition, persistent nocturnal enuresis may adversely affect a child's psychological and social well-being. Children may experience embarrassment, reduced self-esteem, social withdrawal and difficulties in family and school environments. The prevalence of enuresis decreases progressively with increasing age. The synopsis on which this article is based proposes an investigation of the role of individualized homoeopathic treatment in children aged 4–10 years with nocturnal enuresis.

Objective-The primary objective is to assess the effect of homoeopathic medicines in the management of nocturnal enuresis in children aged 4–10 years.

Methods-A case-series study is proposed to be conducted in the OPD/IPD attached to a postgraduate homoeopathic medical college. Thirty cases of nocturnal enuresis will be included using simple random sampling. Children aged 4–10 years of either sex and from all socioeconomic classes will be considered. Cases with gross urological pathology, major systemic disease, malignancy or immunocompromised status will be excluded. Improvement will be assessed using the Enuresis Frequency-Severity Scale, and data will be analyzed using a paired test along with descriptive measures including mean, standard deviation, frequency and percentage.

Conclusion-The proposed study is

intended to generate clinical data regarding the management of nocturnal enuresis with individualized homoeopathic treatment. Its findings may contribute to the understanding of treatment outcomes in pediatric patients; however, conclusions regarding efficacy should be based on the results obtained after completion of the study.

Keywords-Nocturnal enuresis, bedwetting, children, pediatric age group, homoeopathy, individualized treatment, urinary incontinence.

Introduction-Nocturnal enuresis is characterized by repeated involuntary urination during sleep in a child who has reached the developmental stage at which bladder control is expected. The condition is particularly relevant in the pediatric age group because persistent bed-wetting can influence emotional development, self-confidence and social participation.

The synopsis identifies nocturnal enuresis as a prevalent pediatric condition affecting children between 4 and 10 years and emphasizes its possible psychosocial consequences, including embarrassment, social isolation and diminished self-esteem.

The prevalence of enuresis varies with age. The literature reviewed in the synopsis reports estimates of approximately 5–10% among five-year-old children and 1.5–5% among children aged 9–10 years, with prevalence declining further during adolescence.

Nocturnal enuresis is therefore not merely a problem of involuntary urination. For some children and families, recurrent bed-wetting can become a source of anxiety, embarrassment and interpersonal conflict. Relapses may further affect self-image, self-esteem and participation in

social activities.

Homoeopathy approaches patients through individualization of treatment and assessment of the totality of symptoms. The proposed study seeks to evaluate the clinical response of children with nocturnal enuresis receiving homoeopathic treatment.

Definition-The term enuresis is derived from the Greek word enourein, meaning “to void urine.” The term has subsequently become associated particularly with urinary events occurring during sleep. Enuresis may occur during the night, during the day, or during both periods.

For the present study, the focus is on nocturnal enuresis in children aged 4–10 years.

Epidemiology-Enuresis is relatively common during childhood, with prevalence decreasing as children become older. The literature included in the synopsis describes regular enuretic behavior in a considerable proportion of younger children and a progressive decline with age. By approximately 10 years of age, the reported prevalence is considerably lower than in younger children.

The synopsis also reports that nocturnal enuresis is more common in boys and constitutes the majority of childhood enuresis cases. Spontaneous resolution may occur over time, with an estimated annual resolution rate of approximately 15%.

Etiology and Contributing Factors-The development of nocturnal enuresis is multifactorial. The literature reviewed in the synopsis describes contributions from neurobiological mechanisms, bladder function, hormonal regulation, genetic predisposition and psychosocial factors.

Neurobiological factors-Urinary control

involves complex interactions between cerebral and spinal centers, sensory and motor pathways and autonomic and voluntary nervous systems. Regulation of bladder activity involves neural centers in the pons and midbrain.

Genetic factors-A familial tendency has been reported, with higher prevalence among first-degree relatives. Greater concordance among monozygotic twins compared with dizygotic twins suggests an important genetic contribution.

Hormonal and bladder factors-Children with enuresis may have altered patterns of nighttime urine production. Lower nighttime antidiuretic hormone levels have been proposed as one factor contributing to increased nocturnal urine output.

Psychosocial factors-Psychosocial events may precipitate or aggravate enuresis in susceptible children. Examples mentioned in the synopsis include the birth of a sibling, hospitalization, starting school, family separation and moving to a new environment.

Classification-

The synopsis describes three categories:

- Nocturnal-only enuresis – urinary episodes occurring during nighttime sleep.
- Diurnal-only enuresis – urinary episodes occurring during waking hours.
- Nocturnal and diurnal enuresis – urinary episodes occurring during both nighttime and daytime.

The present study focuses specifically on nocturnal enuresis.

Diagnosis and Assessment-Diagnosis requires appropriate clinical assessment and exclusion of medical conditions that may produce urinary symptoms. According to the literature included in the synopsis, diagnostic criteria require an appropriate developmental or chronological age and

recurrent episodes associated with distress or impairment.

A Careful History Should Include:

- Age of onset
- Frequency of bed-wetting
- Previous period of dryness
- Daytime urinary symptoms
- Urgency and frequency
- Bowel habits
- Sleep pattern
- Family history
- Psychosocial circumstances
- Previous treatment
- Associated medical conditions

Laboratory and other investigations are used selectively to exclude organic causes. Urinary tract infection, structural abnormalities, neurological disorders and conditions causing polyuria should be considered in the differential diagnosis.

Differential Diagnosis-

Conditions that should be considered during evaluation include:

- Urinary tract infection
- Bladder dysfunction
- Obstructive uropathy
- Neurological disorders such as spina bifida occulta
- Cystitis
- Diabetes mellitus
- Diabetes insipidus
- Seizure-related urinary incontinence
- Sleepwalking disorder
- Medication-related enuresis

The presence of both nocturnal and diurnal symptoms, particularly when associated with urinary frequency and urgency, warrants careful evaluation for an underlying organic disorder.

Psychosocial Impact-Although nocturnal enuresis may resolve spontaneously, persistent or recurrent episodes can have an important psychosocial impact. The

synopsis describes possible consequences including poor self-image, decreased self-esteem, embarrassment, family conflict and restriction of social activities.

Consequently, management should not focus solely on the urinary symptom. The child's emotional state, family environment and associated behavioral or developmental concerns should also be considered.

Homoeopathic Perspective-The synopsis describes homoeopathic management as being based on individualization and the totality of symptoms. The homoeopathic literature reviewed in the study identifies several medicines traditionally associated with enuresis.

•**Sepia**-Sepia is described in the synopsis in relation to involuntary urination during sleep, particularly when associated with a sensation of weakness or sinking in the abdomen. Associated mental and emotional characteristics described include irritability, indifference and aversion to family members.

•**Causticum**-Causticum is described for involuntary urination during sleep with weakness of bladder sphincter control. Urinary dribbling, urgency and certain forms of urinary incontinence are included among its reported indications in the source literature.

•**Equisetum**-Equisetum is associated in the synopsis with frequent urination during both day and night, bladder fullness and an irritable bladder. Nocturnal enuresis and urinary dribbling are among the urinary symptoms described.

•**Kreosotum**-Kreosotum is described in relation to involuntary urination during the first part of sleep. Strong-smelling urine and difficulty awakening from sleep are also included in the cited materia medica

description.

•**Natrummuriaticum**-Natrummuriaticum is described for involuntary urination, particularly during the first part of sleep, with associated weakness or emotional symptoms such as suppression or grief.

•**Cina**-Cina is described in the synopsis in relation to involuntary urination during sleep, restless sleep and grinding of the teeth, with possible associated irritability and abdominal symptoms.

Selection of an individual medicine should be based on the clinical picture and totality of symptoms rather than solely on the diagnosis of nocturnal enuresis.

Aim and Objectives

•**Aim**-To assess the efficacy of homoeopathic medicines in the management of nocturnal enuresis in children aged 4–10 years.

•**Primary Objective**-To assess the effect of homoeopathic medicines in the management of nocturnal enuresis.

Materials and Methods:

•**Study Design**-The proposed study is a case-series study.

•**Study Setting**-The study is proposed to be conducted in the OPD/IPD attached to a postgraduate homoeopathic medical college.

•**Study Population**-Children diagnosed with nocturnal enuresis constitute the study population.

•**Sample Size**-A total of 30 cases are proposed for inclusion.

•**Sampling Technique**-Simple random sampling is proposed.

Inclusion Criteria-The study includes:

- Children aged 4–10 years.
- Children of either sex.
- Children from all socioeconomic classes.
- Cases diagnosed with the relevant urinary disorder.

Exclusion Criteria-

The following are excluded:

- Children below 4 years or above 10 years.
- Cases with gross urological pathology.
- Cases with major systemic disease.
- Cases with malignancy.
- Immunocompromised cases.

Withdrawal Criteria-Patients who do not

complete follow-up or who deteriorate and require acute life support are to be withdrawn from the study.

Assessment of Treatment Response-The proposed study uses the Enuresis Frequency-Severity Scale as the method of measurement.

Treatment response is categorized as:

•**Improved**: Relief of signs and symptoms up to the end of the study period.

•**Partially improved**: Partial relief of complaints.

•**Not improved**: No change in signs and symptoms.

Data collection includes observation, interview, questionnaire and case study methods, with secondary information obtained from standard textbooks, reference books, journals and research articles.

Data Analysis-The collected data are proposed to be managed in a computerized spreadsheet and presented using appropriate charts and diagrams. The synopsis proposes analysis using a paired test, with results expressed through mean, standard deviation, frequency and percentage.

The statistical analysis will therefore permit comparison of the measured enuresis status before and after the intervention within the study population.

Discussion-Nocturnal enuresis is a common pediatric condition with a multifactorial background. The condition generally becomes less prevalent with increasing age, but persistent symptoms can produce substantial emotional and social consequences.

The literature review included in the synopsis highlights an important evidence gap. A review of complementary and miscellaneous interventions identified 24 randomized or quasi-randomized trials involving 2,334 children, but the quality of many trials was considered poor. Importantly, the review reported that no trials involving homeopathy were included.

This provides a rationale for further systematic investigation. The proposed

study attempts to address this gap by prospectively observing children with nocturnal enuresis and assessing clinical changes following homoeopathic management.

At the same time, because the proposed study is a 30-case case series without a stated control group, any improvement observed cannot by itself establish that homoeopathic treatment caused the improvement. Natural spontaneous resolution, behavioral changes and other factors may influence outcomes. Therefore, the results should be interpreted within the methodological limitations of the study design.

Limitations-The synopsis does not provide completed study results; therefore, treatment efficacy cannot yet be concluded.

Important methodological limitations of the proposed design include:

- Small sample size of 30 cases.
- Case-series design.
- Absence of a stated control or placebo group.
- Potential influence of spontaneous remission.
- Potential influence of behavioral and environmental factors.
- Limited generalizability beyond the selected pediatric population.
- The study duration and final outcome data are not presented in the synopsis.

These limitations should be considered when interpreting the eventual findings

Ethical Considerations-The synopsis includes an informed consent process. Participants or their guardians are informed regarding homoeopathic treatment, possible additional investigations, photography for diagnostic or treatment purposes, alternative treatment options and responsibilities associated with participation. The consent form specifically states that no guarantee or assurance regarding treatment results is provided.

Appropriate evaluation and referral should be undertaken when an underlying medical condition is suspected.

Conclusion-Nocturnal enuresis is a

common pediatric urinary problem that may have significant psychosocial consequences despite its frequently self-limiting nature. Its multifactorial etiology requires appropriate clinical evaluation and exclusion of underlying organic conditions.

The proposed study investigates the role of homoeopathic medicines in children aged 4–10 years using a 30-case case-series design and an Enuresis Frequency-Severity Scale for assessment.

The homoeopathic literature included in the synopsis describes medicines such as Sepia, Causticum, Equisetum, Kreosotum, Natrummuriaticum and Cina in relation to different clinical presentations of enuresis.

However, the present synopsis contains a research-question inconsistency: the primary research question mentions Crotalus horridus, whereas the stated null and alternative hypotheses refer to Cina. This should be corrected in the final dissertation/article before submission so that the research question, hypothesis, intervention and methodology are internally consistent.

Since the uploaded document is a synopsis rather than a completed study report, definitive claims regarding the efficacy of homoeopathic treatment should be reserved until the clinical data, statistical analysis and results are available.

References-The article is based on the literature and references contained in the uploaded synopsis, including works cited from the International Children's Continence Society, Cochrane literature, psychiatric texts, materia medica and repertory sources.

Author-Dr. Qureshi Rukhsar Rajik
MD Part 2 (Paediatric)
Foster Development Homoeopathic
Medical College, Chhatrapati
Sambhajnagar
Under The Guidance of **-Dr. Jagdish Welkar**
(Guide and Professor)