



PRO LIFE ERA
HOMEOPATHY JOURNAL



Dr. Dnyaneshwar
Balasaheb Bhosale

To Study The Homoeopathic Approach In Cases of Pica

Abstract-Pica is a feeding and eating disorder characterised by persistent consumption of non-nutritive, non-food substances for at least one month. Common substances include soil, clay, chalk, ice, paper, starch, ash and other non-food materials. Pica may occur in children, pregnant women and individuals with developmental, psychiatric or nutritional conditions. It may be associated with iron deficiency and anaemia and can cause gastrointestinal, toxicological and dental complications.

Homoeopathy approaches illness through individualisation and considers the totality of characteristic symptoms, including mental, emotional, physical-general and particular symptoms. In cases of pica, detailed case-taking may help identify the patient's cravings, associated mental and physical symptoms, modalities, causative factors, constitutional characteristics and maintaining factors. Repertorisation may be used as an aid to remedy selection.

The purpose of this article is to discuss the possible methodology of a homoeopathic approach in cases of pica. Appropriate medical evaluation remains essential because pica can be associated with nutritional deficiency and serious complications.

Keywords-Pica, Homoeopathy, Constitutional treatment, Craving, Individualisation, Repertorisation, Iron deficiency, Eating disorder.

Introduction-Pica is characterised by persistent consumption of substances that have little or no nutritional value. According to DSM-5 criteria, the behaviour persists for at least one month, is inappropriate for the individual's developmental level and is not culturally or socially sanctioned. Examples include soil, clay, chalk, ice, paper, starch, charcoal, ash and other non-food materials.

Pica can occur at different ages and is particularly observed in children and during pregnancy. It may also occur with intellectual disability and some psychiatric disorders. Nutritional deficiencies, particularly iron

deficiency, are important clinical associations.

From the homoeopathic perspective, pica may be studied not merely as a diagnostic label but as an individual symptom within the patient's totality. The specific substance craved, circumstances in which the craving occurs and associated mental and physical characteristics may be considered during case analysis.

Aim-To study the homoeopathic approach in cases of pica.

Objectives

- To study the clinical presentation of pica.
- To understand associated nutritional and psychological factors.
- To study characteristic symptoms associated with pica from a homoeopathic perspective.
- To evaluate individualisation and constitutional assessment.
- To study the role of repertorisation in remedy selection.
- To assess clinical changes during follow-up.
- To identify the importance of medical investigation and referral.

Definition of Pica-Pica is defined as persistent eating of non-nutritive, non-food substances for at least one month in a manner that is developmentally inappropriate and not culturally accepted.

Common examples include soil or clay (geophagia), ice (pagophagia), raw starch (amylophagia), chalk, paper, ash, charcoal, soap, hair, cloth and other non-food substances.

Association with Iron Deficiency-One of the most important clinical associations of pica is iron deficiency. A 2023 scoping review examining pica and iron-deficiency anaemia identified relevant published evidence and reported that treatment of iron deficiency was associated with resolution of pica symptoms in reported cases.

Depending on the clinical situation, assessment may include complete blood count, haemoglobin, serum ferritin and iron studies, and blood lead testing when exposure is possible. Other investigations should be guided by the substance ingested and clinical presentation.

Homoeopathic Approach to Pica-Individualisation is an important principle of classical homoeopathic case management. The diagnosis of pica alone does not determine the homoeopathic prescription. The physician attempts to understand the complete individual picture.

Important areas of case-taking include:

- Nature and specific substance of the craving
- Frequency, intensity and duration
- Circumstances that increase or decrease the craving
- Mental and emotional characteristics
- Appetite, thirst and food preferences
- Sleep and dreams
- Thermal preferences
- Associated physical complaints
- Causative factors
- Personal and family history
- Nutritional and medical history

Mental symptoms may include anxiety, irritability, fear, restlessness, sadness, anger, grief, emotional sensitivity, attention difficulties and compulsive tendencies. Physical generals include appetite, thirst, sleep, perspiration, bowel habits, energy and other characteristic features.

Repertorial Approach-After complete case-taking, characteristic symptoms may be converted into appropriate repertorial rubrics according to the repertory being used. Possible symptom areas include cravings for specific non-food substances, abnormal appetite and associated mental or physical characteristics.

Repertorisation should not be based solely on the diagnostic term "pica." Greater emphasis may be given to clearly characteristic and individualizing symptoms. The final prescription should be based on the totality of the case and appropriate clinical judgement.

Constitutional Perspective-The constitutional approach attempts to understand the patient as an individual rather than treating an isolated symptom.

Pica symptom → Associated mental state → Physical generals → Causative factors → Associated complaints → Constitutional totality → Repertorisation → Remedy selection → Follow-up

This framework allows the unusual eating behaviour to be interpreted within the broader clinical picture.

Role of Follow-Up-Follow-up is essential for evaluating progress. The following may be monitored:

- Frequency of pica episodes.
- Intensity of craving.
- Amount of substance consumed.
- Ability to resist the craving.
- Associated physical symptoms.
- Mental and emotional symptoms.

- Appetite and sleep.
- General wellbeing.
- Laboratory parameters when abnormal at baseline.
- Occurrence of complications.

Safety and Conventional Medical Assessment-Pica should not be approached exclusively as a homoeopathic symptom because it may indicate nutritional, developmental, behavioural or psychiatric problems.

Medical assessment is particularly important when there is pregnancy, suspected iron deficiency or anaemia, possible lead exposure, abdominal pain, vomiting, suspected intestinal obstruction, ingestion of a toxic substance, severe nutritional deficiency or significant developmental or psychiatric comorbidity.

Identified nutritional deficiencies should be treated using appropriate evidence-based medical care, and behavioural or psychiatric assessment should be considered when clinically indicated.

Discussion-Pica is a clinically significant condition with multiple possible contributing factors. The relationship between pica and iron deficiency is particularly important.

From a homoeopathic perspective, the unusual craving can be considered a potentially characteristic symptom, but the diagnosis alone should not determine treatment. The totality of symptoms, including mental, physical-general and particular symptoms, may be analysed during individualised case-taking.

However, robust clinical evidence demonstrating that homoeopathic medicines independently treat pica is currently insufficient. Therefore, claims of efficacy should be based on properly designed clinical research rather than anecdotal improvement alone. Future studies should use clear diagnostic criteria, adequate sample sizes, appropriate outcome measures, nutritional assessment and transparent reporting of concomitant treatments.

Limitations

- Pica has heterogeneous clinical presentations.
- Underlying causes may differ between individuals.
- Nutritional deficiencies can confound clinical outcomes.
- Spontaneous changes in pica behaviour may occur.

- Concomitant nutritional, behavioural or medical treatment may influence outcomes.
- Evidence specifically evaluating homoeopathic treatment for pica is limited.
- Larger, well-designed prospective clinical studies are required.

Conclusion-Pica is a clinically important disorder characterised by persistent consumption of non-nutritive substances. It may be associated with iron deficiency, nutritional abnormalities, developmental conditions, pregnancy and psychological or behavioural factors.

The homoeopathic approach may involve detailed individualized case-taking, assessment of the specific craving, evaluation of mental and physical generals, identification of causative factors, repertorisation and constitutional analysis.

At the same time, pica requires appropriate medical evaluation because nutritional deficiencies and complications may have significant health consequences. Further high-quality research is necessary to establish the effectiveness and safety of homoeopathy specifically in patients with pica.

References-1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders: DSM-5-TR. 5th ed., text revision. Washington, DC: American Psychiatric Association; 2022.

2. Al Nasser Y, Muco E, Alsaad AJ. Pica. StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; updated 2023.

3. Young SL. Pica in pregnancy: new ideas about an old condition. Annual Review of Nutrition. 2010;30:403-422.

4. Miao D, Young SL, Golden CD. A meta-analysis of pica and micronutrient status. American Journal of Human Biology. 2015.

5. The Association Between Pica and Iron-Deficiency Anemia: A Scoping Review. PubMed-indexed review; 2023.

6. A systematic review of the effectiveness of behavioural treatments for pica in youths. Clinical Psychology Review. 2020.

Dr. Dnyaneshwar Balasaheb Bhosale

MD Part 2 (Paediatrics)

Foster Development Homoeopathic Medical College, Chh Sambhajinagar

Under The Guidance of -

Dr. R.K. Bonde

(Guide and Professor)